



INTERNATIONAL PRELIMINARY EXAMINING

To:

LEE, Sang-Jin

3rd Fl., Dukwon Bldg. 637-19 Yeoksam-dong, Gangnam-gu, 135-080 Seoul, Republic of Korea

NOTIFICATION OF TRANSMITTAL OF INTERNATIONAL PRELIMINARY **EXAMINATION REPORT** 

(PCT Rule 71.1)

Date of mailing

(day/month/year)

01 FEBRUARY 2005 (01.02.2005)

Applicant's or agent's file reference

PCT/KR2003/002692

International application No.

PCT482

IMPORTANT NOTIFICATION

International filing date (day/month/year) 09 DECEMBER 2003 (09.12.2003) Priority date (day/months/year)

12 DECEMBER 2002 (12.12.2002)

Applicant

LEE, Sang Moon

- 1. The applicant is hereby notified that International Preliminary Examining Authority transmits here with the international preliminary examination report and its annexes, if any, established on the international application.
- A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
- Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report(but not of any annexes) and will transmit such translation to those Offices.

#### REMINDER

The applicant must enter the national phase before each elected office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details in the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/KR

Korean Intellectual Property Office 920 Dunsan-dong, Seo-gu, Daejeon 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

COMMISSIONER:

Telephone No. 82-42-481-5131



#### **COPY FOR IB**

## PATENT COOPERATION TREATY

# **PCT**

REC'L	0 1	MAR, 2005
WIPO		PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Artcle 36 and Rule 70)

08 JUN 2005

Applicant's or agent's file reference				
PCT482 FOR FURTHE	SeeNotificationofTransmittalofInternationalPreliminary Examination Report (Form PCT/IPEA/416)			
	g date(day/month/year) Priority date (day/month/year)			
International Patent Classification (IPC) or national classifi	ER 2003 (09.12.2003) 12 DECEMBER 2002 (12.12.2002)			
IPC7 A61B 5/04 Applicant				
LEE, Sang Moon				
<ol> <li>This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</li> <li>This REPORT consists of a total of3sheets, including this cover sheet.</li> <li>This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</li> </ol>				
These annexes consist of a total ofsheets.				
This report contains indications relating to the following items:  I				
Date of submission of the demamd	Date of completion of this report			
12 JULY 2004 (12.07.2004)	31 JANUARY 2005 (31.01.2005)			
Name and mailing address of the IPEA/KR	Authorized officer			
Korean Intellectual Property Office 920 Dunsan-dong, Seo-gu, Daejeon 302-701, Republic of Korea	JEONG, So Yeon			
Facsimile No. 82-42-472-7140	Telephone No. 82-42-481-5656			



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International aplication No. PCT/KR2003/002692

I. Basis of the report		
1. With regard to the elements of the international application:*		
the international application as originally filed  X the description:		
pages 2 5-6 8 10-12 15 18-21		
pages	, as originally filed	
pages 1-1-1,3-4-1,7-7-1,9-9-1,13-14,16-17-1,22-25-1 , filed with the letter of 1	, filed with the demand	
the claims:	2/07/2004	
pages	as originally filed	
pages	with any statment) under Article 19	
pages, filed with the letter of	, filed with the demand	
the drawings:		
pages		
pages	, as originally filed, filed with the demand	
, filed with the letter of	, filed with the demand	
the sequence listing part of the description:		
pages	, as originally filed	
pages		
2. With regard to the language, all the elements marked above and the second se	is Authority in the language in the language	
the international application was filed, unless otherwise indicated under this item.	is realishly in the language in which	
These elements were available or furnished to this Authority in the following language	English which is	
the language of a translation furnished for the purposes of international search (under l	Rule 23.1(b)).	
the language of publication of the international application (under Pule 49.3/5)		
the language of the translation furnished for the purposes of international preliminary or 55.3).	examination(under Rules 55.2 and/	
<ul> <li>3. With regard to any nucleotide and/or amino acid sequence disclosed in the international preliminary examination was carried out on the basis of the sequence listing: <ul> <li>contained inthe international application in written form.</li> <li>filed together with the international application in computer readable form.</li> <li>furnished subsequently to this Authority in written form.</li> <li>furnished subsequently to this Authority in computer readable form</li> <li>The statement that the subsequently furnished written sequence listing does not international applicationas as filed has been furnished.</li> <li>The statement that the information recorded in computer readable form is identical the been furnished.</li> </ul> </li> </ul>	go beyond the disc losure in the	
4. X The amendments have resulted in the cancellation of:		
the description, pages 1,3,4,7,9,13,14,16,17,22-25		
the claims, Nos.		
the drawings, sheets		
This report has been established as if (some of) the amendments had not been made go beyond the disclosure as filed, as indicated in the Supplemental Box(Rule 70.2(c)).	c, since they have been considered to	
Replacement sheets which have been furnished to the receiving Office in response to an invitate in this opinion as "originally filed." and are not annexed to this report since they do not co and 70.17).	ion under Article 14 are referred to ontain amendments (Rules 70.16	
** Any replacement sheet containing such amendments must be referred to under item I and anne	exed to this report.	
rm PCT/IPEA/409 (Box I)(July 1998)		



## INTERNATIONAL PRELIMINARY EXAMINATION

Intor	natio	nal	anlic	

PCT/KR2003/002692

<b>v</b> .	/. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement			
1.	Statement			
	Novelty (N)	Claims	1-2	YES
	Novelty (N)	Claims	None	NO
	Inventive step (IS)	Claims	1-2	YES
	myentive step (15)	Claims	None	NO
	Industrial applicability (IA)	Claims	1-2	YES
	madstrat approaching (= -)	Claims	None	NO
12	Citations and explanations (I	Rule 70.7)		

Claims 1-2 meet the criteria set out in PCT Article 33(2)-(4), because:

(a)the prior art dose not teach or fairly suggest a solid bio-material for a sensor that detects bioelectric signals through the use of the characteristics and functions of bio-epidermal tissues and epidermal tissues of living organisms and the methods for producing the same as claimed in the independent claims of Claims 1 and 2.

CT/KR03/002692 [PEA/KR 12.07.2004

SOLID BIO-MATERIAL FOR A SENSOR THAT DETECTS BIOELECTRIC SIGNALS THROUGH THE USE OF THE
CHARACTERISTICS AND FUNCTIONS OF BIO-EPIDERMAL TISSUES
AND EPIDERMAL TISSUES OF LIVING ORGANISMS AND THE
METHODS FOR PRODUCING THE SAME

### TECHNICAL FIELD

5

10


15

20

The present invention discloses a solid bio-material for the detection of a bio-electromagnetic signal, which senses an information signal generated from living organisms and changes thereof by using the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface having the function of detecting, memorizing and transferring a weak information signal (bio-signal) of an electromagnetic field generated from bio tissues, and a method for producing the same.

#### BACKGROUND ART


AMENDED SHEET (ART. 34)



We have known about electromagnetic signals of bio-organisms since 1 BC. According to documents, the people of that period first tried to cure migraine headaches and prolapse of the anus through the use of electric fish. However, it had not been recognized as a kind of general vital phenomenon not limited to certain species until 1786 that a German biologist, physician, and anatomist, Luigi Galvani (1737-1798) observed electric potential of bio-organisms through experiments using

10

15



potential caused by contractions of the heart muscle and to record them graphically. He coined the term, electrocardiogram for this process.

Therefore, there have been two different methods thus far for detecting bio-signal in general. The first is passively through heat and humidity of the epidermis after adding weak electricity to the exposed epidermis. The second is actively by detecting weak electricity left on the epidermis with electrode.

5


. 10

15

20

However, we cannot detect the smallest differences in biological signals through either of these classical methods due to environmental changes. In other words, technology, thus far, has not been able to distinguish between the ever-slight differences of biological signals of cancer cells and other disease cells.

While searching for a more efficient material to detect the differences in the biological signals of cancer cells we discovered that the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface react very sensitively to the biological signals.




Now, let's take a closer look at the characteristics and functions of the epidermis.

We have originally believed the epidermis to be only a dead keratin tissue layer because of its lack of blood vessels, nerve endings or lymph channels. However, as technology progressed, the epidermis was found to operate a complex organ of numerous structures (sometimes called the integumentary system) serving vital protective and other functions

10

5

15



against external (mechanical, chemical, and physical) offenses.

The ectoderm forms the whole of the nervous system, the epidermis of the skin, the lining cells of the sebaceous, sudoriferous, and mammary glands, the hairs and nails, the epithelium of the nose and adjacent air sinuses, and that of the cheeks and roof of the mouth. From it also are derived the enamel of the teeth, and the anterior lobe of the hypophysis cerebri, the epithelium of the cornea, conjunctiva, and lacrimal glands, and the neuro-epithelium of the sense organs.


When we take a look at a fish scale without its thin outer layer through a microscope, we can find many melanin crystalloids that have been formed by melanin cells when the epidermis has been formed by the ectoderm. These melanin crystalloids have a very complex structure and their shape is similar to that of Neuroglia Cells, namely, the Astrocyte and the Oligodendrocyte.

10

15

20

The epidermis that has been formed by the ectoderm, such as the keratin layer of the human skin, chitinous substance of insects, the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the



horny layer on crustacea's body surface through treatment, will be able to analyze, synthesize, memorize, learn, transform, transmit, and retransmit the electromagnetic signal spectrum of living organisms. In this way, the epidermis' function is similar to that of the human brain.

The epidermis covers the surface of the living organism with the optical medium, which is a semi-transparent, solid, intermittent multi-layer system made of mainly keratin. This intermittent multi-layer system,

10

5

15

PCAKR03/002692

The non-linear medium, optic characteristic and multilayer system of the epidermis produce oscillation harmonic generation when the living organism like the epidermis is influenced by electromagnetic radiation.

5

10

15


20

The epidermis is very different from the other types of living organisms. The epidermis is the border between the outer and inner environments however; the epidermis is also the medium that connects the two environments together. Thus, we can refer to the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface as "multi-information systems".

The epidermis will not simply allow all types of light spectrum, from ultraviolet rays to near infrared rays, to pass through. The epidermis is very selective in what it will allow to pass through and will generate a drastic change in some areas of the spectrum. This is the result of the resonance of the epidermis.

The epidermis will absorb the radiated energy from the electromagnetic field after being stimulated. However, this is not simple





absorption but a modifying procedure of the spectrum of radiated energy. From a biological point of view, this is of great significance. When the radiated spectrum is transferred to short-wave range, excess energy will be generated within the living organism. In other words, the core energy that enhances the activity of the living organism is as such produced.

10

5

15

epidermal layer is composed of a very solid layer system. And, in view of electrophysics, the epidermis is almost same as an insulator having  $10^{12}\sim10^{15}$  ohm. However, the epidermis belongs to the bio-electret under certain conditions. In this case, the bio-electret is the dielectric substance with quasiconstant electric charges. All types of electret charges are very stable. We can observe this electret effect in the biopolymer. The epidermis, which is a biopolymer, optic multi-layer medium, and keratin thin layer, is a natural bio-electret that reacts by generating electric oscillation from the stimulation of an outer electromagnetic field.

5

10

15

The epidermis is a non-linear medium. Since the dielectric constant of non-linear medium is sensitive to the electromagnetic field, the epidermis exposed to a very strong electromagnetic field can generate polarization. Furthermore, this bio-electret can retain polarization even after being removed from the source of polarization.

Even piezoelectricity has been observed on the epidermis. Its characteristics are to be authentic ferroelectrics so that the epidermis is considered "semi-stable ferroelectrics".

The ferroelectrics exposed to the very strong electromagnetic field
has a different non-linear relationship of polarization according to the
intensity of the outer electromagnetic field; in consequence, harmonic
generation is produced in the electric current that is passed through the
dielectric like the epidermis.


AMENDED SHEET (ART. 34)

The epidermal thin multilayer contains keratinocyte generated

5

10

15



Rectification. The wave of polarization of frequency wave 2 will be reradiated at the same frequency wave under the proper conditions. In other words, the second optic harmonic generation of the frequency wave of 2 is produced in the medium.

The square non-linear medium like the epidermis concentrates the wave frequency spectrum when the light wave disperses on the inside. And when the two waves react reciprocally on the standard frequency , the re-radiated wave of the frequency wave 2 is generated. This is the second optic harmonics generation.

5

10

15

20

This is how the non-linear optic medium in the form of the crystalloid produces the second harmonics generation. The epidermis is a crystalloid including a melanin corpuscle.

The epidermis has a Periodic System after which the two layers next to each other can be distinguished by the different physical characteristics like the dielectric and this difference is transferred from one layer to the other several times.

When we examine the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or


scales which contains cuticle, and the horny layer on crustacea's body surface through the microscope, we can observe hundreds of epidermal layers juxtaposed in the micron unit of periodicity.

The two epidermises next to each other in the periodic system have different optico-physical characteristics. First, the refractive values n, n', to two different dielectric constant rates  $\epsilon, \epsilon'$  are different. The non-

10

5

15



linear way the optic harmonic generation is produced is also a characteristic of the periodic medium.

The original characteristic of the periodic medium is that the condition of synchronization is modified when harmonic generation is produced. The diffraction of the harmonic generation derived from the non-linear periodic medium emphasizes the non-linear optic modifying effect. In other words, the periodic medium like the epidermis has optimal conditions for phase synchronization and harmonic generation.

5

10


15

20

The incidence angle range of the bundle of scattered light is wide enough for all types of angles for synchronization. In other words, a general type of lighting can generate phase synchronization even on the epidermal multi-layer structure of the living organism.

The sun energy in range of infrared reaching the earth can influence all types of living organisms while some of this sun energy is reflected on the epidermis. Sometimes, the sun energy is refracted, scattered and finally absorbed on the border of the dielectric layer. The very active non-linear optic medium like the epidermis modifies and shortens the outer energy/infrared light inside of the epidermis.

When multiple monochromatic waves are diffused on the epidermis, its non-linearity generates a combined frequency. The amplitude of each combining wave defines its amplitude. Even if even one of the combining frequencies stays in the visible diapason, the combined frequency will equally remain there. Because only the first combining



signal of the living organism.

## DISCLOSURE OF THE INVENTION

As we have just discussed, our invention is the manufacturing of a solid bio-material with the characteristics of the epidermis by separating the epidermal tissue from a living organism. The characteristics of the epidermis allow us to detect cancer cells by distinguishing signals between cancer cells and normal healthy cells. The characteristics of the epidermis also allow us to produce a more efficient fertilizer. Thus, the purpose of our invention is to create biomaterials for the use in the production of a bio-sensor and the process of manufacturing the bio-sensor.

15


20

5

10

# BEST MODES FOR CARRYING OUT THE INVENTION

The object of our invention is a solid bio-material for the detection of a bio-electromagnetic signal by using epidermal tissues of living organisms by the method of: immersing the carcass of the living organism with the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a



reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface, etc. in a mixed solution of aromatics (fragrance), salt and water; separating the epidermis from the immersed living organism; washing the separated epidermis, soaking it in a mixed solution of potassium dichromate,

10

5

15

vinegar and water, and then drying it in room temperature; applying hot and cold air in turn to the dried epidermis, sterilizing the hot and cold treated epidermis by irradiating ultraviolet rays; generating static electricity; selecting areas with concentrated melanin crystalloid; cutting the epidermis into required sizes and applying pine nut oil to the outer surface. We then take these solid bio-materials for a sensor that detects bio-electric signals.

5

10

15

20

Let's take a detailed look at the methods for producing the solid bio-material for the detection of a bio-electromagnetic signal by using epidermal tissues of living organisms by: immersing the carcass of the living organism with the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface, etc. in a mixed solution of aromatics (fragrance), salt and water in the ratio of 1:2:300 for one week; the water temperature should range between 25-27°C.

The reason for the immersion is to accelerate the decomposition of the carcass that will allow for an easier separation of the epidermis and

AMENDED SHEET (ART. 34)

for the minimalization of scarring on the epidermis.

For maximum melanin crystalloid, the epidermis should be thick and semi-transparent. Semi-transparency is necessary for the optical function of the epidermis while the thickness will allow for an easier creation of harmonic generation energy.

The separating procedure is as follows: the immersed the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface should be separated from the decomposed living organism by

5

10

- 1. The external electromagnetic spectrum can transform the epidermis, which is close to an insulator having electric resistance of  $10^{12}\sim10^{15}\Omega$  to a Dielectric Substance conducting polarization.
- 2. The keratinocyte of the epidermis causes non-linear electromagnetic polarization under the influence of electromagnetic spectrum.
  - 3. The epidermis is the bio-electret, which can retain polarization even without the external primary source of polarization.
- 4. The bio-electret that has been influenced by the electromagnetic

  field produces Harmonic Generation due to the changes of the nonlinear relationship of polarization.


As shown above, the optical, physical and electrical characteristics of the micro multilayer epidermal becomes dramatically active through our biosensor manufacturing procedure after separating from the organism.

15

20

The followings examples show specific application of the solid bio materials for the detection of bio-signal using epidermal tissues of living organism according to the present invention may be applied to the following fields.

First, in the case of the reaction of the biosensor to energy of living organisms, our solid living bio-material can detect, analyze, and synthesize the electromagnetic spectrum of living organisms that have



abnormal cells such as cancer or tumor cells. So, if we set up this solid living bio-material on the head of the probe, the solid living bio-material will detect and amplify even the smallest capacitance from the living organism. Our probe is equipped with three special circuits; the first circuit enables the probe to amplify and transform this capacitance into frequency waves; the second enables the probe to take the frequency waves and diagnose them; the third enables the probe to transform the diagnoses and translate it into digital form, which is displayed on an LCD panel.


Second, in the case of the fertilizing aspect of the solid living biomaterial, it can rejuvenate soil and aid in the growth of crops and plants. For this purpose, we utilized bird feathers. The solid living biomaterial treated through our procedure can revive uncultivable soil by stimulating the solid living organism in soil and enhancing growth of crops and plants.

10

15

In order to test for the fertilization purpose of our solid living biomaterial, we sowed barley seeds in 1) soil mixed with invented solid living bio-material and 2) soil mixed with non-treated bird feather.


Note: All conditions were held constant between two soil tests and results were taken three days after disseminating seeds.



Unit: cm

	The growth rate of barley grown in soil mixed with		
	invented solid living	non-treated bird feathers	
	organism sample (treated)		
3 days after			
disseminating	0.2	<del>-</del> .	
seeds in soil	·		
4 days	4	1	
5 days	6	2.4	
б days	10.1	5 .	
7 days	14	7	
8 days	18	9.5	
9 days	20	13.5	
10 days	20	14	
11 days	19.5	14	
12 days	21	14	
13 days	21.75	14.5	
14 days	23	15	
15 days	24	16	
i6 days	24.5	17	
17 days	25.5	17.5	
18 days	26	18	

AMENDED SHEET (ART. 34)



19 days	27	18.5
20 days	27.5	19.5
21 days	29	20.5
22 days	30	21
23 days	30	22
24 days	32	22.5
25 days	33	24

According to the above results, we found that barley seeds strewn in soil mixed with invented solid living organism grew at a much faster rate than barley seed strewn in soil mixed with non-treated bird feathers.


5

10

15

#### INDUSTRIAL APPLICABILITY

This invention gives us the new biological materials in order to enhance human life. The biological materials may be used for detecting cancer or creating a revolutionary fertilizer. The new materials are made of the epidermis of living organisms, scales which have been generated from dermis, as well as the deformation of skin which came from a degeneration or keratinization of scales, fish scales, the scales or horny substances of a reptile(tortoises)'s body surface area, the deformed body surface skin of birds and mammals, the feathers of birds(fowl), the body surface of insects, mollusca, shellfish and the cuticle on their body



surfaces, vertabrata feathers or scales which contains cuticle, and the horny layer on crustacea's body surface which can detect, analyze, synthesize, memorize, transform and transmit electric charges from the skin.

5

10

15

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.